

Original Article

EVALUATION OF ANTIOXIDANT, ANTICHOLINESTERASE AND ANTIPROLIFERATIVE ACTIVITIES OF THE AERIAL PARTS OF *DRYOPTERIS RADDEANA*

Asuman Evrim Uygun¹ , Hüsniye Aka Sağlıker¹ , Mustafa Sevindik*¹ 

¹ Department of Biology, Faculty of Engineering and Nature Sciences, Osmaniye Korkut Ata University, Osmaniye, 80000, Turkey.

* Correspondence, e-mail: sevindik27@gmail.com

Received: 21.05.2025 / Revised: 17.07.2025 / Accepted: 06.08.2025 / Published online: 20.08.2025 /
Published in final form: 21.12.2025

ABSTRACT

This study aimed to evaluate the biological activities of the ethanol extract obtained from the aerial parts of *Dryopteris raddeana* (Fomin) Fomin. Total antioxidant capacity (TAS), total oxidant level (TOS) and oxidative stress index (OSI) values determined by Rel Assay kits using the ethanol extract obtained from the aerial parts of the plant were determined as 5.099 ± 0.076 mmol/L, 7.354 ± 0.107 μ mol/L, and 0.144 ± 0.004 , respectively. These results show that *D. raddeana* exhibits a balanced oxidative profile and has a low oxidant load. In anticholinesterase activity analyses, the inhibitory effect of the extract against acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) enzymes was determined as 58.99 ± 1.48 μ g/mL and 83.51 ± 1.59 μ g/mL, respectively. In the evaluation of the antiproliferative effect, it was observed that extract concentrations in the range of 25–200 μ g/mL applied to the A549 human lung cancer cell line decreased cell viability in a dose-dependent manner. A significant decrease in cell viability was detected especially at concentrations of 100 and 200 μ g/mL. In conclusion, in this study, it was determined that the plant could be an antioxidant, antimicrobial, and antiproliferative source for these biological activities of *D. raddeana*.

KEYWORDS: *Dryopteris raddeana*, antioxidant activity, anticholinesterase, antiproliferative effect, phytotherapy.

Article is published under the CC BY license.

1. Introduction

Plants have been at the forefront of natural resources that support human health throughout history; they have been widely used in traditional treatment systems with their medicinal, aromatic, and nutritional properties [1]. Today, plants have become the focus of modern pharmaceutical research thanks to the wide biological activities of the secondary metabolites they contain [2]. Phytochemicals such as alkaloids, flavonoids, phenolic compounds, terpenoids, tannins, and saponins exhibit a wide range of biological effects, including antioxidant, antimicrobial, anticancer, anti-inflammatory, hepatoprotective, and antidiabetic activities. [3–9]. In recent years, interest in natural products has increased due to the side effects and resistance problems of synthetic drugs. In this context, determining the biological activities of extracts obtained from various plant species is of great importance both in terms of the discovery of new drug candidates and the development of natural treatment alternatives [10,11]. In addition, performing biological activity tests at in vitro and in vivo levels scientifically

reveals the pharmacological potential of these extracts and contributes to the support of traditional knowledge with modern methods.

Dryopteris raddeana (Fomin) Fomin (old name: *Dryopteris pallida*) is a fern species belonging to the Pteridaceae family, widespread in high mountainous regions. This species is known for growing in moist and shady habitats and has attracted attention in systematic botanical studies with its morphological differences [12]. Recent studies have revealed that ferns are important not only in terms of taxonomy but also in terms of their chemical contents and potential biological activities. In this context, the chemical components of *Dryopteris raddeana*, especially phloroglucide derivatives, have been studied in detail by advanced semi-quantitative analysis methods. Phloroglucides are phenolic compounds generally known for their antioxidant, antimicrobial, and cytotoxic effects [13]. The high levels of these compounds in *D. raddeana* were found to be remarkable in terms of the potential pharmacological effects of the species. In addition, it is thought that the secondary metabolites contained in this species may

have protective effects in biological systems related to oxidative stress [14]. In this context, advanced biological activity tests on *D. raddeana* may contribute to new natural product-based treatment approaches. In this study, it was aimed to determine the antioxidant, antiproliferative, and anticholinesterase activities of *D. raddeana*.

2. Materials and Methods

The *Dryopteris raddeana* samples used in this study were obtained from Fenk Yaylası, Osmaniye Province, Turkey. The aerial parts (leaves and stems) of the plant were dried in a laboratory environment at room temperature (-22 – 24°C), in the shade, for 7 days to preserve phytochemical stability. During the extraction process, 10 grams of plant material were taken and extracted in a Soxhlet apparatus for approximately 6 hours with 250 mL of 96% ethanol at 50°C . The obtained crude extract was separated from its solvent using a Buchi R100 rotary evaporator device operating at 40°C . The resulting extracts were stored at $+4^{\circ}\text{C}$ until experimental applications. All measurements were performed in triplicate, and results are presented as mean \pm standard deviation (StD). No statistical significance tests were applied.

2.1. Antioxidant tests

In this study, the total antioxidant capacity (TAS) and total oxidant level (TOS) of the extract obtained from the aerial parts of *Dryopteris raddeana* using ethanol were determined using commercial analysis kits belonging to the Rel Assay company. The measurements were carried out in accordance with the analysis protocols provided by the manufacturer. While trolox was used as the standard substance in TAS analyses, calibration was performed with hydrogen peroxide for TOS analyses. TAS results were reported in mmol/L, and TOS results in $\mu\text{mol/L}$ [15,16]. In the calculation of the oxidative stress index (OSI), both parameters were converted to the same unit, and the ratio obtained by dividing TOS by TAS was presented in percentage format. OSI was calculated using the formula: $\text{OSI (arbitrary unit)} = \text{TOS } [\mu\text{mol H}_2\text{O}_2 \text{ equivalent/L}] / (\text{TAS [mmol Trolox equivalent/L]} \times 10)$ [17].

2.2. Anticholinesterase tests

In this study, anticholinesterase activities of extracts of the aerial parts of *D. raddeana* were evaluated based on the colorimetric method developed by Ellman et al. [18]. In inhibitory activity assays against acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) enzymes, galantamine was used as the standard inhibitor. Stock solutions were prepared by diluting plant extracts in the concentration range of 200 to $3.125 \mu\text{g/mL}$. In the experimental procedure, $130 \mu\text{L}$ (0.1 M , $\text{pH } 8.0$) phosphate buffer, $10 \mu\text{L}$ of test sample, and $20 \mu\text{L}$ of enzyme solution (AChE or BChE) were added to each microwell, respectively; the mixture was incubated at 25°C in the dark for 10 min. After incubation, $20 \mu\text{L}$ of DTNB [5,5'-dithiobis(2-nitrobenzoic acid)] solution, and then $20 \mu\text{L}$ of substrate (acetylcholine iodide or butyrylcholine iodide) were added to initiate the reaction. Enzymatic activity was measured spectrophotometrically at 412 nm wavelength. Each test was performed in triplicate, and the inhibition percentages of the extracts were calculated and expressed as IC_{50} ($\mu\text{g/mL}$) values.

2.3. Antiproliferative activity tests

In this study, the cytotoxic effect of the aerial parts of *D. raddeana* extract on the A549 human lung carcinoma cell line was evaluated using the MTT (methyl thiazolyl tetrazolium) cell viability test method. Before experimental applications, plant extracts were prepared at concentrations of 25, 50, 100, and $200 \mu\text{g/mL}$ to obtain stock solutions. After the cells reached approximately 70–80% density, the surface detachment was performed with 3.0 mL Trypsin-EDTA solution (Sigma-Aldrich, MO, USA). The separated cells were transferred to culture plates at an appropriate density and incubated for 24 hours. Following incubation, plant extracts prepared at different doses were applied to the cells, and a second 24-hour incubation period was started. At the end of the process, the supernatant medium was removed, and MTT solution at a concentration of 1 mg/mL was added, and the cells were incubated at 37°C . After the formation of formazan crystals was observed, dimethyl sulfoxide (DMSO) (Sigma-Aldrich, MO, USA) was added to dissolve these crystals. Cell viability was measured at a wavelength of 570 nm using an Epoch brand spectrophotometer (BioTek Instruments, Winooski, VT, USA) [19].

3. Results and Discussion

3.4. Antioxidant Activity

Plants are exposed to various environmental stress factors throughout their growth and development, and these stresses can result in excessive production of reactive oxygen species (ROS), leading to oxidative stress [20]. The resulting oxidant compounds may cause lipid peroxidation, protein denaturation, and DNA damage in plant cells. In order to prevent such molecular damage, plants activate their antioxidant systems as a natural defense strategy [21,22]. These antioxidant components protect cell integrity by neutralizing the harmful effects of ROS and increase the plant's resistance to stress conditions. Therefore, determining plant antioxidant capacity is essential both for understanding the adaptation mechanisms against environmental stresses and for biotechnological and pharmacological applications [23,24]. In this study, the antioxidant potential of the aerial parts of *D. raddeana* was evaluated. The obtained data are presented in Table 1.

Table 1. TAS, TOS ve OSI values of *Dryopteris raddeana*.

Plant	TAS mmol/L	TOS $\mu\text{mol/L}$	OSI
<i>Dryopteris raddeana</i>	5.099 ± 0.076	7.354 ± 0.107	0.144 ± 0.004

* Values are presented as mean \pm SD

There is no direct study on the antioxidant activity of *D. raddeana* in the literature. However, there are various scientific findings on the antioxidant potential of different *Dryopteris* species [25–27]. In this context, our study was carried out to evaluate the antioxidant properties of *D. raddeana*. In our study, the total antioxidant capacity (TAS), total oxidant level (TOS) and oxidative stress index (OSI) values of the plant were measured using Rel Assay kits. Data on these parameters of different plant species have been widely reported in the literature. For comparison, TAS values of *Scorzonera*

papposa, *Mentha longifolia*, *Anthemis cotula*, *Hypericum spectabile*, *Equisetum ramosissimum* and *Ferulago platycarpa* were determined as 6.328, 6.094, 7.625, 9.306, 4.802, and 5.688 mmol/L, respectively. TOS values were 11.525, 14.050, 11.247, 13.065, 7.643, and 15.552 μ mol/L, respectively; while OSI values were reported as 0.182, 0.231, 0.148, 0.140, 0.159, and 0.273 [28–33].

Compared to these studies, the TAS value of *D. raddeana* was higher than that of *E. ramosissimum* and lower than those of *S. papposa*, *M. longifolia*, *A. cotula*, *H. spectabile*, and *F. platycarpa*. TAS is an important indicator reflecting the total antioxidant compounds in a natural product [34]; therefore *D. raddeana* can be considered to possess a moderate antioxidant capacity. Especially when compared to some species with lower TAS values, it is understood that *D. raddeana* has a remarkable potential in terms of antioxidant compound content. The TOS value represents the total oxidant load in a natural product [34]. In our study, the TOS value of *D. raddeana* was found to be lower than that of all comparison species, indicating that the plant contains low levels of oxidant compounds and its potential to create oxidative stress is limited. In this respect, *D. raddeana* exhibits a positive profile, especially in terms of antioxidant/oxidant balance. In addition, this low oxidant load suggests that the plant may be a safe candidate for phytotherapeutic uses. The OSI value is a critical parameter reflecting oxidative stress balance. A low OSI indicates that antioxidants exhibit effective protection against oxidant compounds [34]. The OSI value of *D. raddeana* was found to be higher than only that of *H. spectabile* and lower than that of all other comparison species. This indicates that the plant's antioxidant defense system effectively combats oxidative load.

Overall, the antioxidant profile of *D. raddeana*, supported by its low oxidant load and balanced OSI value, indicates its potential as a valuable candidate for the discovery of new natural antioxidant sources. In addition, more detailed determination of the potential antioxidant compounds of this species with different extraction methods and fractionation studies may expand its pharmacological applications.

3.5. Anticholinesterase activity

In recent years, studies on the anticholinergic activities of plant-derived natural compounds have gained momentum. Inhibition of acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) enzymes has become an important pharmacological target, especially in the treatment of neurodegenerative disorders such as Alzheimer's disease [35]. Many medicinal and aromatic plant species have the potential to inhibit these enzymes through the alkaloids, flavonoids, terpenoids, and phenolic compounds they contain. These compounds obtained from plant extracts can support synaptic transmission by increasing acetylcholine levels and slowing down the progression of cognitive disorders. In this context, the discovery of natural products with anticholinergic effects is of great importance both in terms of developing safe alternative treatment strategies and in revealing new drug candidates for the pharmaceutical industry [36,37]. In this study, the anticholinesterase activity of *D. raddeana* was investigated, and the results are presented in Table 2.

Table 2. Anti-AChE and anti-BChE values of *Dryopteris raddeana*.

Sample	AChE μ g/mL	BChE μ g/mL
<i>Dryopteris raddeana</i>	58.99 \pm 1.48	83.51 \pm 1.59


* Values are presented as mean \pm SD.

Literature reviews revealed no previous studies on the anticholinesterase activity of *D. raddeana*. However, several data exist regarding the inhibitory effects of other species of the same genus on acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) enzymes [25,38]. This underscores the potential contribution of our study evaluating the anticholinesterase properties of *D. raddeana*. In our study, the inhibitory activity of the aerial parts of *D. raddeana* against AChE and BChE enzymes was evaluated using galantamine as the reference standard. It was found that galantamine showed high levels of inhibition on both enzymes. This result, as expected, is consistent with galantamine being a clinically approved anticholinesterase agent. On the other hand, it was found that *D. raddeana* extract also showed a certain level of inhibitory effect on both enzymes. This finding indicates that the plant structure may contain natural compounds with anticholinesterase potential (e.g. phenolic compounds, flavonoids, or terpenoids). However, the fact that the inhibitory activity is not as strong as galantamine may be due to the complex structure of the herbal extract and the presence of active compounds in diluted concentrations.

Anticholinesterase activity is an important treatment target, especially in neurodegenerative disorders such as Alzheimer's disease [39]. Therefore, determining the AChE and BChE inhibitory capacity of *D. raddeana* is important for the preliminary evaluation of the pharmacological potential of the species. In addition, the obtained data may pave the way for the isolation of more specific and potent inhibitory compounds by biodirected fractionation studies of this plant. In conclusion, these first findings regarding the anticholinesterase activity of *D. raddeana* may form the basis for future in vivo and molecular level studies. Particularly, purified extracts or the association of specific phytochemical compounds with biomarker activities will enable the evaluation of this species as a potential natural resource in neuroprotective drug development studies.

3.6. Antiproliferative activity

Plants are among the important natural resources in anticancer drug research thanks to their rich content of secondary metabolites. It has been scientifically proven that compounds such as flavonoids, alkaloids, terpenoids, phenolic acids, and lignans, which are among these metabolites, have antiproliferative effects on many cancer cell lines [40,41]. Antiproliferative activity is usually achieved by mechanisms such as arresting the cell cycle at certain stages, triggering apoptotic pathways, or mitochondrial dysfunction. These effects of plant extracts often occur both in a dose-dependent manner and in a cell type-specific manner. In this respect, plant sources offer valuable alternatives in terms of both efficacy and toxicity in the development of chemotherapeutic drugs [42,43]. In this study, the effects of *D. raddeana* on A549 lung cancer cells were investigated, and the results are presented in Figure 1.



(Control: The group not treated with chemicals, only kept in the medium; DMSO: The group in which the medium and DMSO were applied; the group in which the extract was applied at 25, 50, 100 and 200 µg/mL concentrations, OD: Optical Density measured at 570 nm using MTT assay)

Fig. 2. Antiproliferative effect of *D. raddeana* aerial part extract on A549 cells

In this study, the antiproliferative effects of *D. raddeana* ethanol extract on the A549 human lung adenocarcinoma cell line was evaluated, and the obtained findings are presented in Figure 1. Cell viability decreased significantly depending on the concentration of the applied extract. The significant decrease observed especially at 100 and 200 µg/mL concentrations reveals the dose-dependent cytotoxic effect of the extract. According to the data in Figure 1, no decrease in cell viability was observed in the DMSO group compared to the control group. This shows that DMSO did not show a toxic effect under experimental conditions, and the observed cytotoxic effect was specific to *D. raddeana* extract. The decrease in OD values in the extract-applied groups reveals that cell proliferation was suppressed, and the extract showed antiproliferative effects. Possible mechanisms of this effect may include induction of apoptosis, arrest of the cell cycle, or increased cellular damage via oxidative stress.

The absence of any findings in the literature regarding the effects of *D. raddeana* on the A549 cell line increases the originality and scientific contribution of this study. However, it has been reported that some *Dryopteris* species belonging to the same genus exhibit antiproliferative effects against A549 cells [44–46]. These previous studies show that *Dryopteris* species are rich in secondary metabolites and that these compounds may have anticancer properties. In this context, it is thought that *D. raddeana* may also contain similar structural components. The findings obtained reveal that *D. raddeana* may be a potential anticancer agent by reducing the viability of A549 cells at certain concentrations. The importance of plant sources is increasing, especially in natural product research, and revealing the potential of species such as *D. raddeana* in this area may contribute to new therapeutic strategies. In future studies, biodirected fractionation of the extract, isolation of active compounds, and elaboration of the mechanistic effects of these compounds at in vitro and in vivo levels are of great importance. In addition, analyses at the gene expression level will help to explain which molecular pathways this species affects to produce antiproliferative effects.

4. Conclusions

This study is the first comprehensive investigation evaluating the antioxidant, anticholinesterase, and antiproliferative potential of the aerial parts of *D. raddeana*. The results showed that the total antioxidant capacity (TAS) of *D. raddeana* was significant, while its total oxidant level (TOS) was low; therefore, the oxidative stress index (OSI) had a balanced profile. These findings indicate that the plant has a strong antioxidant profile and is a natural source that can be evaluated for pharmaceutical applications.

In addition, in the anticholinesterase analyses, it was determined that *D. raddeana* extract showed a certain level of inhibitory effect against both acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) enzymes. It is thought that these effects may be related to the phenolic or flavonoid compounds in the structure of the plant. Although it did not exhibit as high an inhibitory activity as galantamine, these results indicate that the plant can be evaluated among natural inhibitors for neurodegenerative diseases. In antiproliferative activity tests, it was determined that *D. raddeana* extract reduced cell viability in a dose-dependent manner on the A549 lung cancer cell line. The significant decrease observed especially at 100 and 200 µg/mL concentrations indicates that the plant can be considered as a potential anticancer agent. The absence of any study on this species in the literature specifically for the A549 cell line makes this study original and provides a new contribution to the literature.

In conclusion, the antioxidant, anticholinesterase, and antiproliferative activities of *D. raddeana* indicate that this species should be considered among natural biological agents. Future studies should further explore these findings through biodirected fractionation of the plant extract, compound isolation, and molecular level mechanism analyses. Thus, it can be demonstrated that *D. raddeana* may be a potential phytotherapeutic source for pharmaceutical, neuroprotective, and oncological applications.

Author Contributions: Conceptualization, A.E.U., H.S., and M.S.; methodology, A.E.U., H.S., and M.S.; validation, A.E.U., H.S., and M.S.; investigation, A.E.U., H.S., and M.S.; resources, A.E.U., H.S., and M.S.; data curation, A.E.U., H.S., and M.S.; writing-original draft preparation, A.E.U., H.S., and M.S.; writing-review and editing, A.E.U., H.S., and M.S.; visualization, A.E.U., H.S., and M.S.; supervision, A.E.U., H.S., and M.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Mohammed, F.S.; Uysal, I.; Sevindik, M. A review on antiviral plants effective against different virus types. *Prospects Pharm. Sci.* **2023**, *21*(2), 1-21. DOI: 10.56782/pps.128
2. Sevindik, M.; Mohammed, F.S.; Uysal, I. Autism: plants with neuro-psychopharmacotherapeutic potential.

- Prospects Pharm. Sci. 2023, 21(3), 38-48. DOI: 10.56782/pps.143
3. Turkmen, D.; Dursun, A.H.; Caliskan, O.; Koksak Kavrak, M.; Guler, Z. Volatile compounds, phenolic content, and antioxidant capacity in Sultan hawthorn (*Crataegus azarolus* L.) leaves. *J. Agric. Sci. Technol.* 2023, 25(5), 1089-1099. DOI: 10.22034/jast.25.5.1089
4. Martínez-Olivo, A.O.; Zamora-Gasga, V.M.; Medina-Torres, L.; Pérez-Larios, A.; Sáyo-Ayerdi, S.G.; Sánchez-Burgos, J.A. Biofunctionalization of natural extracts, trends in biological activity and kinetic release. *Adv. Colloid Interface Sci.* 2023, 318, Art. No: 102938. DOI: 10.1016/j.cis.2023.102938
5. Uysal, I.; Koçer, O.; Mohammed, F.S.; Lekesiz, Ö.; Doğan, M.; Şabik, A.E.; Sevindik, E.; Gerçekler, F.Ö.; Sevindik, M. Pharmacological and nutritional properties: Genus *Salvia*. *Adv. Pharmacol. Pharm.* 2023, 11, 140-155. DOI: 10.13189/app.2023.110205
6. Dursun, A.; Çalışkan, O.; Güler, Z.; Bayazit, S.; Türkmen, D.; Gündüz, K. Effect of harvest maturity on volatile compounds profiling and eating quality of hawthorn (*Crataegus azarolus* L.) fruit. *Sci. Hortic.* 2021, 288, Art. No: 110398. DOI: 10.1016/j.scienta.2021.110398
7. Koçer, O. Determination of optimum extract conditions and evaluation of biological activity potential of *Salvia cilicica* Boiss. *Sci. Rep.* 2025, 15(1), Art. No: 9277. DOI: 10.1038/s41598-025-93925-2
8. Yazar, M.; Sevindik, M.; Uysal, I.; Polat, A.O. Effects of caffeic acid on human health: Pharmacological and therapeutic effects, biological activity and toxicity. *Pharm. Chem. J.* 2025, 59, 49-55. DOI: 10.1007/s11094-025-03363-7
9. Colak, S.; Comlekcioglu, N.; Aygan, A.; Kocabas, Y.Z.; Comlekcioglu, U. Phytochemical properties and bioactive potential of various *Astragalus* spp. from Turkey. *Food Biosci.* 2025, 64, Art. No: 105901. DOI: 10.1016/j.fbio.2025.105901
10. Koçer, O.; Pham, T.V.; Yaz, H.H.; Kaya, H.; Açıkgöz, A. Antioxidant and anticholinesterase activities of *Stachys pumila*: Biological activities of *Stachys pumila*. *Eurasian J. Med. Biol. Sci.* 2024, 4(2), 80-84.
11. Yazar, M.; Sevindik, M.; Polat, A.O.; Koçer, O.; Karatepe, H.K.; Uysal, İ. General properties, biosynthesis, pharmacological properties, biological activities and daily uses of luteolin. *Prospects Pharm. Sci.* 2024, 22(4), 146-154. DOI: 10.56782/pps.286
12. Güner, A.; Kandemir, A. (Eds.). Resimli Türkiye Florası: Cilt 2; Ali Nihat Gökyiğit Vakfı Yayınları: İstanbul, Türkiye, 2018.
13. Nardi, E. La distribuzione italiana di «*Dryopteris pallida*» (Bory) Fomin. *Webbia* 1976, 30, 3-32. DOI: 10.1080/00837792.1976.10670042
14. Widen, C.J.; Vida, G.; Von Euw, J.; Reichstein, T.V. Die Phlorglucide von *Dryopteris viliarii* (Bell.) Woynar und anderer Farne der Gattung *Dryopteris* sowie die mögliche Abstammung von *D. filix-mas* (L.) Schott. *Helv. Chim. Acta* 1971, 54, 2824-2850. DOI: 10.1002/hlca.19710540850
15. Erel, O. A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. *Clin. Biochem.* 2004, 37(4), 277-285. DOI: 10.1016/j.clinbiochem.2003.11.015
16. Erel, O. A new automated colorimetric method for measuring total oxidant status. *Clin. Biochem.* 2005, 38, 1103-1111. DOI: 10.1016/j.clinbiochem.2005.08.008
17. Sevindik, M. Anticancer, antimicrobial, antioxidant and DNA protective potential of mushroom *Leucopaxillus gentianeus* (Qué.) Kotl. *Indian J. Exp. Biol.* 2021, 59, 310-315. DOI: 10.56042/ijeb.v59i05.50501
18. Ellman, G.L.; Courtney, K.D.; Andres, V., Jr.; Featherstone, R.M. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* 1961, 7, 88-95. DOI: 10.1016/0006-2952(61)90145-9
19. Bal, C.; Akgül, H.; Sevindik, M.; Akata, İ.; Yumrutaş, Ö. Determination of the anti-oxidative activities of six mushrooms. *Fresenius Environ. Bull.* 2017, 26, 6246-6252.
20. Çömlekçioğlu, N.; Aygan, A.; Sevindik, M.; Çömlekçioğlu, U. Comparative analysis of phytochemical properties of extracts obtained from flowers and leaves of *Astragalus angustifolius* collected from different locations. *Prospects Pharm. Sci.* 2024, 22(3), 62-68. DOI: 10.56782/pps.214
21. Baba, H.; Sevindik, M.; Doğan, M.; Akgül, H. Antioxidant, antimicrobial activities and heavy metal contents of some Myxomycetes. *Fresenius Environ. Bull.* 2020, 29, 7840-7846.
22. Eraslan, E.C.; Altuntaş, D.; Baba, H.; Bal, C.; Akgül, H.; Akata, İ.; Sevindik, M. Some biological activities and element contents of ethanol extract of wild edible mushroom *Morchella esculenta*. *Sigma J. Eng. Nat. Sci.* 2021, 39, 24-28. DOI: 10.14744/sigma.2021.00003
23. Elbozan, H.; Korkmaz, N. An examination of the antioxidant and antibacterial capabilities, as well as the phenolic compounds, of *Silene macrodonta* and *Silene chaetodonta*. *Prospects Pharm. Sci.* 2025, 23(2), 35-41. DOI: 10.56782/pps.287
24. Akkaya, O.B.; Çelik, İ.S.; Ertaş, E.; Çömlekçioğlu, N.; Aygan, A. Determination of antimicrobial, anticarcinogenic activity of bioactive components of *Hypericum perforatum* L. plant. *Int. J. Chem. Technol.* 2024, 8, 73-82. DOI: 10.32571/ijct.1445857
25. Alam, F.; Khan, S.H.A.; Asad, M.H.H.B. Phytochemical, antimicrobial, antioxidant and enzyme inhibitory potential of medicinal plant *Dryopteris ramosa* (Hope) C. Chr. *BMC Complement. Med. Ther.* 2021, 21, Art. No: 10. DOI: 10.1186/s12906-021-03370-7
26. Rani, A.; Uzair, M.; Ali, S.; Qamar, M.; Ahmad, N.; Abbas, M.W.; Esatbeyoglu, T. *Dryopteris juxtaopatia* root and shoot: Determination of phytochemicals; antioxidant, anti-inflammatory, and hepatoprotective effects; and toxicity assessment. *Antioxidants* 2022, 11, Art. No: 1670. DOI: 10.3390/antiox11091670
27. Akpotu, A.E.; Ghasi, S.I.; Ajibo, D.N.; Ike, A.O.; Akhator, A.J.; Enya, J.I.; Onyeleonu, I. Free radical scavenging activity and phytochemical investigation of ethanol leaf extract of *Dryopteris dilatata*. *J.*

- Complement. Altern. Med. Res. **2023**, 22, 37-48. DOI: 10.9734/jocamr/2023/v22i2489
28. Mohammed, F.S.; Günel, S.; Şabik, A.E.; Akgül, H.; Sevindik, M. Antioxidant and antimicrobial activity of *Scorzonera papposa* collected from Iraq and Turkey. *KSU J. Agric. Nat.* **2020**, 23, 1114-1118. DOI: 10.18016/ksutarimdogan.vi.641944
 29. Mohammed, F.S.; Günel, S.; Pehlivan, M.; Doğan, M.; Sevindik, M.; Akgül, H. Phenolic content, antioxidant and antimicrobial potential of endemic *Ferulago platycarpa*. *Gazi Univ. J. Sci.* **2020**, 33, 670-677. DOI: 10.35378/gujs.707555
 30. Gürgeç, A.; Sevindik, M.; Krupodorova, T.; Uysal, İ.; Ünal, O. Biological activities of *Hypericum spectabile* extract optimized using artificial neural network combined with genetic algorithm application. *BMC Biotechnol.* **2024**, 24, Art. No: 83. DOI: 10.1186/s12896-024-00914-w
 31. Korkmaz, N.; Koçer, O.; Fathi, S.; Uysal, İ.; Sevindik, M. Branched horsetail (*Equisetum ramosissimum*): some biological activities and total phenolic and flavonoid contents. *Prospects Pharm. Sci.* **2024**, 22(3), 69-75. DOI: 10.56782/pps.209
 32. Sabik, A.E.; Sevindik, M.; Mohammed, F.S.; Doğan, M. A new natural source against A549 lung cancer cells: *Anthemis cotula* and its biological activities and phenolic contents. *Pharm. Chem. J.* **2024**, 58, 1134-1140. DOI: 10.1007/s11094-024-03098-5
 33. Sevindik, M.; Gürgeç, A.; Krupodorova, T.; Uysal, İ.; Koçer, O. A hybrid artificial neural network and multi-objective genetic algorithm approach to optimize extraction conditions of *Mentha longifolia* and biological activities. *Sci. Rep.* **2024**, 14, Art. No: 31403. DOI: 10.1038/s41598-024-31403-0
 34. Gürgeç, A.; Sevindik, M. Application of artificial neural network coupling multiobjective particle swarm optimization algorithm to optimize *Pleurotus ostreatus* extraction parameters. *J. Food Process. Preserv.* **2022**, 46, Art. No: e16949. DOI: 10.1111/jfpp.16949
 35. Korkmaz, N.; Mohammed, F.S.; Uysal, İ.; Sevindik, M. Antioxidant, antimicrobial and anticholinesterase activity of *Dittrichia graveolens*. *Prospects Pharm. Sci.* **2023**, 21(4), 48-53. DOI: 10.56782/pps.169
 36. Şahin, H.; Demir, S.; Boğa, M.; Sarı, A.; Makbul, S.; Gültepe, M. Comparative phytochemical and bioactivity studies on two related *Scorzonera* L. species: A chemotaxonomic contribution. *Biochem. Syst. Ecol.* **2023**, 111, Art. No: 104743. DOI: 10.1016/j.bse.2023.104743
 37. Ersoy, E.; Boğa, M.; Kaplan, A.; Mataraci Kara, E.; Eroglu Ozkan, E.; Demirci Kayiran, S. LC-HRMS profiling of phytochemicals with assessment of antioxidant, anticholinesterase, and antimicrobial potentials of *Astragalus brachystachys* DC. *Chem. Biodivers.* **2025**, 22, Art. No: e202401853. DOI: 10.1002/cbdv.202401853
 38. Cao, J.; Xia, X.; Chen, X.; Xiao, J.; Wang, Q. Characterization of flavonoids from *Dryopteris erythrosora* and evaluation of their antioxidant, anticancer and acetylcholinesterase inhibition activities. *Food Chem. Toxicol.* **2013**, 51, 242-250. DOI: 10.1016/j.fct.2012.09.015
 39. Korkmaz, N. Extract optimization of *Ulva lactuca* L. and biological activities of optimized extracts. *BMC Biotechnol.* **2025**, 25(1), 21. DOI: 10.1186/s12896-025-00954-w
 40. Keskin, C.; Aktepe, N.; Yükselten, Y.; Sunguroglu, A.; Boğa, M. In-vitro antioxidant, cytotoxic, cholinesterase inhibitory activities and anti-genotoxic effects of *Hypericum retusum* Aucher flowers, fruits and seeds methanol extracts in human mononuclear leukocytes. *Iran. J. Pharm. Res.* **2017**, 16, 210. DOI: 10.22037/ijpr.2017.2111
 41. Oran, S.A.; Yousef, I.; Jaffal, S.M. Evaluation of the antiproliferative effect of selected plant extracts on colon and skin cancer cell lines. *J. Appl. Pharm. Sci.* **2022**, 12, 91-96. DOI: 10.7324/JAPS.2022.121110
 42. Özkan, E.E.; Boğa, M.; Yılmaz, M.A.; Kara, E.M.; Yeşil, Y. LC-MS/MS analyses of *Ziziphora clinopodioides* Lam. from Turkey: Antioxidant, anticholinesterase, antimicrobial and anticancer activities. *Istanbul J. Pharm.* **2020**, 50, 33-41. DOI: 10.26650/IstanbulJPharm.2020.0061
 43. Nawaz, A.; Jamal, A.; Arif, A.; Parveen, Z. In vitro cytotoxic potential of *Solanum nigrum* against human cancer cell lines. *Saudi J. Biol. Sci.* **2021**, 28, 4786-4792. DOI: 10.1016/j.sjbs.2021.04.062
 44. Liu, Z.D.; Zhao, D.D.; Jiang, S.; Xue, B.; Zhang, Y.L.; Yan, X.F. Anticancer phenolics from *Dryopteris fragrans* (L.) Schott. *Molecules* **2018**, 23, Art. No: 680. DOI: 10.3390/molecules23030680
 45. Jana, F.; Pareek, S.; Srivastava, R.P.; Zahoor, I.; Sharma, A.; Shrivastava, D. Anti-cancerous and antibacterial potential of silver nanoparticles synthesized using leaf extract of fern *Dryopteris barbigera*. *Dig. J. Nanomater. Biostruct.* **2022**, 17, 285-299.
 46. Said, K.; Rauf, M.; Khan, S.A.; Hussain, A.; Alhegaili, A.S.; Hussain, S.; Hamayun, M. Metabolomics and anticancer potential of the aerial parts of *Dryopteris ramosa* against cancerous cell lines assisted with advanced computational approaches. *Curr. Pharm. Des.* **2025**, 31, 797-820. DOI: 10.2174/1381612829666231227103013